arXiv:1603.05716v1 [math.CA] 14 Mar 2016

Approximation properties on q-Szász-Mirakjan-Kantrovich Stancu type operators via Dunkl generalization

M. Mursaleen and **Md. Nasiruzzaman**
Department of Mathematics, Aligarh Muslim University, Aligarh–202002, India
mursaleenm@gmail.com; nasir3489@gmail.com

Abstract

This paper is devoted to study the approximation properties and rate of approximation of the Szász-Mirakjan-Kantrovich-Stancu type polynomials generated by the Dunkl generalization of the exponential function with respect to q-calculus. We present approximation properties with the help of well-known Korovkin's theorem and determine the rate of convergence in terms of classical modulus of continuity, the class of Lipschitz functions, Peetre's K-functional, and the second-order modulus of continuity. Moreover, we obtain the approximation results for Bivariate case for these operators.

Keywords and phrases: q-integers; Dunkl analogue; Szász operator; q- Szász-Mirakjan-Kantrovich; modulus of continuity; Peetre's K-functional.

AMS Subject Classification (2010): 41A25, 41A36, 33C45.

1. Introduction and preliminaries

S.N Bernstein [2] introduced the very well known Bernstein Classical operators in 1912 as

$$B_n(f;x)=\sum_{k=0}^{n}\binom{n}{k}x^k(1-x)^{n-k}f\left(\frac{k}{n}\right),\ x\in[0,1]. \tag{1.1}$$

for $n\in\mathbb{N}$ and $f\in C[0,1]$.

In 1950, for $x\geq 0$, Szász [25] introduced the operators

$$S_n(f;x)=e^{-nx}\sum_{k=0}^{\infty}\frac{(nx)^k}{k!}f\left(\frac{k}{n}\right),\ f\in C[0,\infty). \tag{1.2}$$

The first q-analogue of the well-known Bernstein polynomials was introduced by Lupaş [12], by applying the idea of q-integers and investigated its approximating and shape-preserving properties. In 1997 Phillips [22] considered another q-analogue of the classical Bernstein polynomials. Later on, many authors introduced q-generalizations of various operators and investigated several approximation properties [13, 14, 15, 16, 17] etc. Recently, making use of (p,q)-calculus, the first (p,q)-analogue of Bernstein operators was given in [18]. There is a generalization in (p,q)-calculus.

The q-integer $[n]_q$, the q-factorial $[n]_q!$ and the q-binomial coefficient are defined by (see [1])

$$[n]_q := \begin{cases} \frac{1-q^n}{1-q}, & \text{if } q \in \mathbb{R}^+ \setminus \{1\} \\ n, & \text{if } q = 1, \end{cases} \quad \text{for } n \in \mathbb{N} \text{ and } [0]_q = 0,$$

$$[n]_q! := \begin{cases} [n]_q[n-1]_q \cdots [1]_q, & n \geq 1, \\ 1, & n = 0, \end{cases}$$

$$\left[\begin{array}{c} n \\ k \end{array} \right]_q := \frac{[n]_q!}{[k]_q![n-k]_q!},$$

respectively.
The q-analogue of $(1+x)^n$ is defined by

$$(1+x)_q^n := \begin{cases} (1+x)(1+qx)\cdots(1+q^{n-1}x) & n = 1, 2, 3, \cdots \\ 1 & n = 0. \end{cases}$$

The Gauss binomial formula is given by

$$(x+a)_q^n = \sum_{k=0}^{n} \left[\begin{array}{c} n \\ k \end{array} \right]_q q^{k(k-1)/2} a^k x^{n-k}.$$

In 1930 the first Bernstein-Kantorovich operators [10] in 1930 and in 1950 Száasz operators [25] were defined. In the recent years Sucu [24] define a Dunkl analogue of Szász operators via a generalization of the exponential function given by [23] as follows:

$$S_n^*(f;x) := \frac{1}{e_\mu(nx)} \sum_{k=0}^{\infty} \frac{(nx)^k}{\gamma_\mu(k)} f\left(\frac{k+2\mu\theta_k}{n}\right), \tag{1.3}$$

where $x \geq 0,\ f \in C[0,\infty), \mu \geq 0,\ n \in \mathbb{N}$
and

$$e_\mu(x) = \sum_{n=0}^{\infty} \frac{x^n}{\gamma_\mu(n)}.$$

Here

$$\gamma_\mu(2k) = \frac{2^{2k} k! \Gamma\left(k+\mu+\frac{1}{2}\right)}{\Gamma\left(\mu+\frac{1}{2}\right)},$$

and

$$\gamma_\mu(2k+1) = \frac{2^{2k+1} k! \Gamma\left(k+\mu+\frac{3}{2}\right)}{\Gamma\left(\mu+\frac{1}{2}\right)}.$$

Cheikh et al. [3] stated the q-Dunkl classical q-Hermite type polynomials and gave definitions of q-Dunkl analogues of exponential functions and recursion relations for $\mu > -\frac{1}{2}$ and $0 < q < 1$.

$$e_{\mu,q}(x) = \sum_{n=0}^{\infty} \frac{x^n}{\gamma_{\mu,q}(n)},\ x \in [0,\infty) \tag{1.4}$$

$$E_{\mu,q}(x) = \sum_{n=0}^{\infty} \frac{q^{\frac{n(n-1)}{2}} x^n}{\gamma_{\mu,q}(n)},\ x \in [0,\infty) \tag{1.5}$$

$$\gamma_{\mu,q}(n+1)=\left(\frac{1-q^{2\mu\theta_{n+1}+n+1}}{1-q}\right)\gamma_{\mu,q}(n),\ n\in\mathbb{N}, \tag{1.6}$$

$$\theta_n=\begin{cases}0 & \text{if } n\in 2\mathbb{N},\\ 1 & \text{if } n\in 2\mathbb{N}+1.\end{cases}$$

Recently İçöz and Çekim gave a Dunkl generalization of Kantrovich integral generalization of Szász operators in [8] and a Dunkl generalization of Szász operators via q-calculus in [9].

This paper is dedicated to construct Kantrovich Stancu type Szász-Mirakjan operators generated by the Dunkl generalization of the exponential function. We study approximation properties of the operators via a universal Korovkins type theorem and a weighted Korovkins type theorem and the rate of convergence of the operators for functions belonging to the Lipschitz class are presented.

2. AUXILIARY RESULTS

We define a Dunkl generalization of Szász-Mirakjan-Kantrovich Stancu type operators via q-calculs:

For any $x\in[0,\infty),\ n\in\mathbb{N},\ 0<q<1$, and $\mu>\frac{1}{2}$, we define

$$\tilde{K_{n,q}}(f;x)=\frac{[n]_q}{E_{\mu,q}([n]_q x)}\sum_{k=0}^{\infty}\frac{([n]_q x)^k}{\gamma_{\mu,q}(k)}q^{\frac{k(k-1)}{2}}\int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} f\left(\frac{nt+\alpha}{n+\beta}\right)d_qt, \tag{2.1}$$

where f is continuous and nondecreasing function on $[0,\infty)$. When we take $\alpha=\beta=0$ in the operator $\tilde{K_{n,q}}(f;x)$ (2.1), then we obtain the Kantorovich-type generalization of Szász operators [19].

Lemma 2.1. *Let $K^*_{n,q}(.\ ;\ .)$ be the operators given by (2.1). Then we have:*

(1) $\tilde{K_{n,q}}(1;x)=1$

(2) $\tilde{K_{n,q}}(t;x)=\frac{n}{(n+\beta)}\left(\frac{\alpha}{n}+\frac{1}{[2]_q[n]_q}\right)+\frac{2nq}{[2]_q(n+\beta)}x$

(3) $\frac{n^2}{(n+\beta)^2}\left(\frac{\alpha^2}{n^2}+\frac{2\alpha}{[2]_q n[n]_q}+\frac{1}{[3]_q[n]_q^2(n+\beta)}\right)+\frac{qn^2}{(n+\beta)^2}\left(\frac{3q^{2\mu+1}[1+2\mu]_q}{[3]_q[n]_q}+\frac{3}{[3]_q[n]_q}+\frac{4\alpha}{[2]_q n}\right)x+\frac{3qn^2}{[3]_q(n+\beta)^2}x^2\le\tilde{K_{n,q}}(t^2;x)\le\frac{n^2}{(n+\beta)^2}\left(\frac{\alpha^2}{n^2}+\frac{2\alpha}{[2]_q n[n]_q}+\frac{1}{[3]_q[n]_q^2(n+\beta)}\right)+\frac{n^2}{(n+\beta)^2}\left(\frac{3[1+2\mu]_q}{[3]_q[n]_q}+\frac{3q}{[3]_q[n]_q}+\frac{4q\alpha}{[2]_q n}\right)x+\frac{3n^2}{[3]_q(n+\beta)^2}x^2.$

Lemma 2.2. *Let the operators $\tilde{K_{n,q}}(.\ ;\ .)$ be given by (2.1). Then we have,*

(1) $\tilde{K_{n,q}}(t-1;x)=\frac{n}{(n+\beta)}\left(\frac{\alpha}{n}+\frac{1}{[2]_q[n]_q}-\frac{n+\beta}{n}\right)+\frac{2nq}{[2]_q(n+\beta)}x$

(2) $\tilde{K_{n,q}}(t-x;x)=\frac{n}{(n+\beta)}\left(\frac{\alpha}{n}+\frac{1}{[2]_q[n]_q}\right)+\left(\frac{2nq}{[2]_q(n+\beta)}-1\right)x$

(3) $\tilde{K_{n,q}}((t-x)^2;x)\le\frac{n^2}{(n+\beta)^2}\left(\frac{\alpha^2}{n^2}+\frac{2\alpha}{[2]_q n[n]_q}+\frac{1}{[3]_q[n]_q^2(n+\beta)}\right)+\left\{\frac{n^2}{(n+\beta)^2}\left(\frac{3[1+2\mu]_q}{[3]_q[n]_q}+\frac{3q}{[3]_q[n]_q}+\frac{4q\alpha}{[2]_q n}\right)-\frac{2n}{(n+\beta)}\left(\frac{\alpha}{n}+\frac{1}{[2]_q[n]_q}\right)\right\}x+\left(\frac{3n^2}{[3]_q(n+\beta)^2}-\frac{2nq}{[2]_q(n+\beta)}+1\right)x^2$

3. Korovkin type approximation properties

In this section, we obtain the Korovkin type and weighted Korovkin type approximation properties by the operators defined in (2.1).

Korovkin-type theorems furnish simple and useful tools for ascertaining whether a given sequence of positive linear operators, acting on some function space, is an approximation process.

Theorem 3.1. *([11, 5]) Let $L_n : C[a,b] \to C[a,b]$ be a sequence of positive linear operators. If $\lim_{n\to\infty} L_n t^j = t^j$, $j = 0,1,2$, uniformly on $[a,b]$, then $\lim_{n\to\infty} L_n f = f$ uniformly on $[a,b]$ for every $f \in C[a,b]$.*

Let $C_B(\mathbb{R}^+)$ be the set of all bounded and continuous functions on $\mathbb{R}^+ = [0,\infty)$, which is linear normed space with

$$\| f \|_{C_B} = \sup_{x\geq 0} | f(x) | .$$

Also let

$$H := \{f : x \in [0,\infty), \frac{f(x)}{1+x^2} \text{ is convergent as } x \to \infty\}.$$

In order to obtain the convergence results for the operators $\tilde{K_{n,q}}(.,.)$, we take $q = q_n$ where $q_n \in (0,1)$ and satisfying,

$$\lim_n q_n \to 1, \ \lim_n q_n^n \to a \tag{3.1}$$

Theorem 3.2. *Let $q = q_n$ satisfy (3.1), for $0 < q_n < 1$ and if $\tilde{K_{n,q_n}}(. \ ; \ .)$ be the operators given by (2.1). Then for any function $f \in C[0,\infty) \cap H$,*

$$\tilde{K_{n,q_n}}(f;x) = f(x)$$

is uniformly on each compact subset of $[0,\infty)$.

Proof. The proof is based on the well known Korovkin's theorem regarding the convergence of a sequence of linear positive operators, so it is enough to prove the conditions

$$\tilde{K_{n,q_n}}((t^j;x) = x^j, \ j = 0,1,2, \ \{\text{as } n \to \infty\}$$

uniformly on $[0,1]$.
Clearly from (3.1) and $\frac{1}{[n]_{q_n}} \to 0$ $(n \to \infty)$, we have

$$\lim_{n\to\infty} K^*_{n,q_n}(t;x) = x, \ \lim_{n\to\infty} K^*_{n,q_n}(t^2;x) = x^2.$$

This complete the proof. □

The weighted Korovkin-type theorems were proved by Gadzhiev [6]. A real function $\rho = 1 + x^2$ is called a weight function if it is continuous on $\mathbb{R}$ and

$$\lim_{|x|\to\infty} \rho(x) = \infty, \ \rho(x) \geq 1 \text{ for all } x \in \mathbb{R}$$

We recall the weighted spaces of the functions on $\mathbb{R}^+$, which are defined as follows:

$$\begin{aligned} P_\rho(\mathbb{R}^+) &= \{f :| f(x) | \leq M_f \rho(x)\}, \\ Q_\rho(\mathbb{R}^+) &= \left\{f : f \in P_\rho(\mathbb{R}^+) \cap C[0,\infty)\right\}, \\ Q_\rho^k(\mathbb{R}^+) &= \left\{f : f \in Q_\rho(\mathbb{R}^+) \text{ and } \lim_{x\to\infty} \frac{f(x)}{\rho(x)} = k(k \text{ is a constant})\right\}, \end{aligned}$$

where $\rho(x) = 1 + x^2$ is a weight function and M_f is a constant depending only on f. Note that $Q_\rho(\mathbb{R}^+)$ is a normed space with the norm $\| f \|_\rho = \sup_{x\geq 0} \frac{|f(x)|}{\rho(x)}$.

Theorem 3.3. *Let $q = q_n$ satisfy (3.1) for $0 < q_n < 1$ and let $\tilde{K_{n,q_n}}(.\ ;\ .)$ be the operators given by (2.1). Then for any function $f \in Q_\rho^k(\mathbb{R}^+)$ we have*

$$\lim_{n\to\infty} \| \tilde{K_{n,q_n}}(f;x) - f \|_\rho = 0.$$

Proof. From Lemma 2.1, the first condition of (1) is fulfilled for $\tau = 0$. Now for $\tau = 1, 2$ it is easy to see from (2), (3) of Lemma 2.1 by using (3.1) that

$$\| \tilde{K_{n,q_n}}(t)^\tau; x) - x^\tau \|_\rho = 0.$$

This complete the proof. □

4. Rate of Convergence

Here we calculate the rate of convergence of operators (2.1) by means of modulus of continuity and Lipschitz type maximal functions.

Let $f \in C[0,\infty]$. The modulus of continuity of f denoted by $\omega(f,\delta)$ gives the maximum oscillation of f in any interval of length not exceeding $\delta > 0$ and it is given by the relation

$$\omega(f,\delta) = \sup_{|y-x|\leq\delta} | f(y) - f(x) |,\ x, y \in [0,\infty). \tag{4.1}$$

It is known that $\lim_{\delta\to 0+} \omega(f,\delta) = 0$ for $f \in C[0,\infty)$ and for any $\delta > 0$ one has

$$| f(y) - f(x) | \leq \left(\frac{| y - x |}{\delta} + 1\right) \omega(f,\delta). \tag{4.2}$$

Theorem 4.1. *Let $\tilde{K_{n,q}}(.\ ;\ .)$ be the operators defined by (2.1). Then for $f \in \tilde{C}[0,\infty),\ x \geq 0,\ 0 < q < 1$ we have*

$$| \tilde{K_{n,q}}(f;x) - f(x) | \leq \left(1 + \sqrt{\lambda_n(x)}\right) \omega\left(f; \frac{1}{\sqrt{[n]_q}}\right),$$

where $\lambda_n(x) = \tilde{K_{n,q}}((t-x)^2;x)$ is defined in (3) of Lemma 2.2 and $\tilde{C}[0,\infty)$ is the space of uniformly continuous functions on $\mathbb{R}^+$, furthermore, $\omega(f,\delta)$ is the modulus of continuity of the function $f \in \tilde{C}[0,\infty)$ defined in (4.1).

Proof. We prove it by using (4.1),(4.2) and Cauchy-Schwarz inequality.
$\mid \tilde{K_{n,q}}(f;x) - f(x) \mid$

$$
\begin{aligned}
&\leq \frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \frac{([n]_q x)^k}{\gamma_{\mu,q}(k)} q^{\frac{k(k-1)}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} \mid f(t) - f(x) \mid d_q(t) \\
&\leq \frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \frac{([n]_q x)^k}{\gamma_{\mu,q}(k)} q^{\frac{k(k-1)}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} \left(1 + \frac{1}{\delta} \mid t - x \mid\right) d_q(t) \omega(f;\delta) \\
&= \left\{1 + \frac{1}{\delta} \left(\frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \frac{([n]_q x)^k}{\gamma_{\mu,q}(k)} q^{\frac{k(k-1)}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} \mid t - x \mid d_q(t) \right) \right\} \omega(f;\delta) \\
&\leq \left\{1 + \frac{1}{\delta} \left(\frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \frac{([n]_q x)^k}{\gamma_{\mu,q}(k)} q^{\frac{k(k-1)}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} (t-x)^2 d_q(t) \right)^{\frac{1}{2}} \left(\tilde{K_{n,q}}(1;x) \right)^{\frac{1}{2}} \right\} \\
&\quad \times \omega(f;\delta) \\
&= \left\{1 + \frac{1}{\delta} \left(\tilde{K_{n,q}}(t-x)^2; x \right)^{\frac{1}{2}} \right\} \omega(f;\delta)
\end{aligned}
$$

if we choose $\delta = \delta_n = \sqrt{\frac{1}{[n]_q}}$, then we get our result. □

Now we give the rate of convergence of the operators $K^*_{n,q}(f;x)$ defined in (2.1) in terms of the elements of the usual Lipschitz class $Lip_M(\nu)$.

Let $f \in C[0,\infty)$, $M > 0$ and $0 < \nu \leq 1$. The class $Lip_M(\nu)$ is defined as

$$Lip_M(\nu) = \{f :\mid f(\zeta_1) - f(\zeta_2) \mid \leq M \mid \zeta_1 - \zeta_2 \mid^{\nu} \ \ (\zeta_1, \zeta_2 \in [0,\infty))\}. \tag{4.3}$$

Theorem 4.2. *Let $\tilde{K_{n,q}}(.\ ;\ .)$ be the operators defined in (2.1). Then for each $f \in Lip_M(\nu)$, $(M > 0,\ 0 < \nu \leq 1)$ satisfying (5.4) we have*

$$\mid \tilde{K_{n,q}}(f;x) - f(x) \mid \leq M \left(\lambda_n(x)\right)^{\frac{\nu}{2}}$$

where $\lambda_n(x) = \tilde{K_{n,q}}\left((t-x)^2; x\right)$ defined in Theorem 4.1.

Proof. We prove it by using (5.4) and Hölder's inequality.

$$
\begin{aligned}
\mid \tilde{K_{n,q}}(f;x) - f(x) \mid &\leq \mid \tilde{K_{n,q}}(f(t) - f(x); x) \mid \\
&\leq \tilde{K_{n,q}}\left(\mid f(t) - f(x) \mid; x\right) \\
&\leq \mid M\tilde{K_{n,q}}\left(\mid t - x \mid^{\nu}; x\right).
\end{aligned}
$$

Therefore

$$
\begin{aligned}
&| \tilde{K_{n,q}}(f;x) - f(x) | \\
&\leq M\frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \frac{([n]_q x)^k}{\gamma_{\mu,q}(k)} q^{\frac{k(k-1)}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} | t-x |^{\nu} \, d_q(t) \\
&\leq M\frac{[n]_q}{E_{\mu,q}([n]_q x)} \sum_{k=0}^{\infty} \left(\frac{([n]_q x)^k q^{\frac{k(k-1)}{2}}}{\gamma_{\mu,q}(k)} \right)^{\frac{2-\nu}{2}} \\
&\times \left(\frac{([n]_q x)^k q^{\frac{k(k-1)}{2}}}{\gamma_{\mu,q}(k)} \right)^{\frac{\nu}{2}} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} | t-x |^{\nu} \, d_q(t) \\
&\leq M \left(\frac{[n]_q}{(E_{\mu,q}([n]_q x))} \sum_{k=0}^{\infty} \frac{([n]_q x)^k q^{\frac{k(k-1)}{2}}}{\gamma_{\mu,q}(k)} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} d_q(t) \right)^{\frac{2-\nu}{2}} \\
&\times \left(\frac{[n]_q}{(E_{\mu,q}([n]_q x))} \sum_{k=0}^{\infty} \frac{([n]_q x)^k q^{\frac{k(k-1)}{2}}}{\gamma_{\mu,q}(k)} \int_{\frac{[k+2\mu\theta_k]_q}{q^{k-2}[n]_q}}^{\frac{[k+1+2\mu\theta_k]_q-1}{q^{k-1}[n]_q}+\frac{1}{[n]_q}} | t-x |^2 \, d_q(t) \right)^{\frac{\nu}{2}} \\
&= M \left(\tilde{K_{n,q}}(t-x)^2; x \right)^{\frac{\nu}{2}}.
\end{aligned}
$$

Which complete the proof. □

Let $C_B[0,\infty)$ denote the space of all bounded and continuous functions on $\mathbb{R}^+ = [0,\infty)$ and

$$C_B^2(\mathbb{R}^+) = \{g \in C_B(\mathbb{R}^+) : g', g'' \in C_B(\mathbb{R}^+)\}, \tag{4.4}$$

with the norm

$$\| g \|_{C_B^2(\mathbb{R}^+)} = \| g \|_{C_B(\mathbb{R}^+)} + \| g' \|_{C_B(\mathbb{R}^+)} + \| g'' \|_{C_B(\mathbb{R}^+)}, \tag{4.5}$$

also

$$\| g \|_{C_B(\mathbb{R}^+)} = \sup_{x \in \mathbb{R}^+} | g(x) | . \tag{4.6}$$

Theorem 4.3. *Let $\tilde{K_{n,q}}(.\ ;\ .)$ be the operators defined in (2.1). Then for any $g \in C_B^2(\mathbb{R}^+)$ we have*

$$| \tilde{K_{n,q}}(f;x) - f(x) | \leq \left\{ \frac{n}{(n+\beta)} \left(\frac{\alpha}{n} + \frac{1}{[2]_q [n]_q} \right) + \left(\frac{2nq}{[2]_q (n+\beta)} - 1 \right) x + \frac{\lambda_n(x)}{2} \right\} \| g \|_{C_B^2(\mathbb{R}^+)}$$

where $\lambda_n(x)$ is given in Theorem 4.1.

Proof. Let $g \in C_B^2(\mathbb{R}^+)$. Then by using the generalized mean value theorem in the Taylor series expansion we have

$$g(t) = g(x) + g'(x)(t-x) + g''(\psi)\frac{(t-x)^2}{2}, \ \psi \in (x,t).$$

By linearity, we have

$$\tilde{K_{n,q}}(g,x) - g(x) = g'(x)\tilde{K_{n,q}}\left((t-x);x\right) + \frac{g''(\psi)}{2}\tilde{K_{n,q}}\left((t-x)^2;x\right),$$

which implies that
$| \tilde{K_{n,q}}(g;x) - g(x) |$

$$\leq \left(\tilde{K_{n,q}}\left((t-x);x\right)\right) \| g' \|_{C_B(\mathbb{R}^+)} + \left(\tilde{K_{n,q}}\left((t-x)^2;x\right)\right) \frac{\| g'' \|_{C_B(\mathbb{R}^+)}}{2}.$$

From (4.5) we have $\| g' \|_{C_B[0,\infty)} \leq \| g \|_{C_B^2[0,\infty)}$.
$| \tilde{K_{n,q}}(g;x) - g(x) |$

$$\leq \left(\tilde{K_{n,q}}\left((t-x);x\right)\right) \| g \|_{C_B^2(\mathbb{R}^+)} + \left(\tilde{K_{n,q}}\left((t-x)^2;x\right)\right) \frac{\| g \|_{C_B^2(\mathbb{R}^+)}}{2}.$$

This completes the proof from Lemma 2.2. □

The Peetre's K-functional is defined by

$$K_2(f,\delta) = \inf_{C_B^2(\mathbb{R}^+)} \left\{ \left(\| f - g \|_{C_B(\mathbb{R}^+)} + \delta \| g'' \|_{C_B^2(\mathbb{R}^+)} \right) : g \in \mathcal{W}^2 \right\}, \tag{4.7}$$

where

$$\mathcal{W}^2 = \left\{ g \in C_B(\mathbb{R}^+) : g', g'' \in C_B(\mathbb{R}^+) \right\}. \tag{4.8}$$

Then there exits a positive constant $C > 0$ such that $K_2(f,\delta) \leq C\omega_2(f,\delta^{\frac{1}{2}})$, $\delta > 0$, where the second order modulus of continuity is given by

$$\omega_2(f,\delta^{\frac{1}{2}}) = \sup_{0<h<\delta^{\frac{1}{2}}} \sup_{x\in\mathbb{R}^+} | f(x+2h) - 2f(x+h) + f(x) | . \tag{4.9}$$

Theorem 4.4. *Let $\tilde{K_{n,q}}(.\ ;\ .)$ be the operators defined in (2.1) and $C_B[0,\infty)$ be the space of all bounded and continuous functions on $\mathbb{R}^+$. Then for $x \in \mathbb{R}^+$, $f \in C_B(\mathbb{R}^+)$ we have*
$| \tilde{K_{n,q}}(f;x) - f(x) |$

$$\leq 2M\left\{ \omega_2\left(\sqrt{\frac{\lambda_n(x)}{4} + \frac{n}{2(n+\beta)}\left(\frac{\alpha}{n} + \frac{1}{[2]_q[n]_q}\right) + \frac{1}{2}\left(\frac{2nq}{[2]_q(n+\beta)} - 1\right)x} \right) \right.$$

$$\left. + \min\left(1, \frac{\lambda_n(x)}{4} + \frac{n}{2(n+\beta)}\left(\frac{\alpha}{n} + \frac{1}{[2]_q[n]_q}\right) + \frac{1}{2}\left(\frac{2nq}{[2]_q(n+\beta)} - 1\right)x\right) \right\},$$

where M is a positive constant, $\lambda_n(x)$ is given in Theorem 4.1 and $\omega_2(f;\delta)$ is the second order modulus of continuity of the function f defined in (4.9).

Proof. We prove this by using Theorem (4.3)

$$\begin{aligned} | \tilde{K_{n,q}}(f;x) - f(x) | &\leq | \tilde{K_{n,q}}(f-g;x) | + | \tilde{K_{n,q}}(g;x) - g(x) | + | f(x) - g(x) | \\ &\leq 2 \| f - g \|_{C_B(\mathbb{R}^+)} + \frac{\lambda_n(x)}{2} \| g \|_{C_B^2(\mathbb{R}^+)} \\ &+ \left\{ \frac{n}{(n+\beta)}\left(\frac{\alpha}{n} + \frac{1}{[2]_q[n]_q}\right) + \left(\frac{2nq}{[2]_q(n+\beta)} - 1\right)x \right\} \| g \|_{C_B(\mathbb{R}^+)} \end{aligned}$$

From (4.5) clearly we have $\| g \|_{C_B[0,\infty)} \leq \| g \|_{C_B^2[0,\infty)}$.
Therefore,
$| \tilde{K_{n,q}}(f;x) - f(x) |$

$$\leq 2 \| f-g \|_{C_B(\mathbb{R}^+)} + \left\{ \frac{\lambda_n(x)}{2} + \frac{n}{(n+\beta)}\left(\frac{\alpha}{n} + \frac{1}{[2]_q[n]_q}\right) + \left(\frac{2nq}{[2]_q(n+\beta)} - 1\right)x \right\} \| g \|_{C_B^2(\mathbb{R}^+)}$$

where $\lambda_n(x)$ is given in Theorem 4.1.

By taking infimum over all $g \in C_B^2(\mathbb{R}^+)$ and by using (4.7), we get

$$| \tilde{K_{n,q}}(f;x) - f(x) | \leq 2K_2 \left\{ f; \frac{\lambda_n(x)}{4} + \frac{n}{2(n+\beta)} \left(\frac{\alpha}{n} + \frac{1}{[2]_q [n]_q} \right) + \frac{1}{2} \left(\frac{2nq}{[2]_q (n+\beta)} - 1 \right) x \right\}$$

Now for an absolute constant $C > 0$ in [4] we use the relation

$$K_2(f;\delta) \leq C\{\omega_2(f;\sqrt{\delta}) + \min(1,\delta) \parallel f \parallel\}.$$

This completes the proof of the theorem. □

5. Construction of Bivariate Operators

In this section, we construct a bivariate extension of the operators (2.1).

Let $\mathbb{R}_+^2 = [0,\infty) \times [0,\infty)$, $f : C(\mathbb{R}_+^2) \to \mathbb{R}$ and $0 < q_{n_1}, q_{n_2} < 1$. We define the bivariate extension of the Dunkl q-parametric Szász-Mirakjan operators (2.1) as follows:

$$K^*_{n_1,n_2}(f;q_{n_1},q_{n_2};x,y) = \frac{1}{E_{\mu_1,q_{n_1}}([n_1]_{q_{n_1}}x)} \frac{1}{E_{\mu_2,q_{n_2}}([n_2]_{q_{n_2}}y)} \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \frac{([n_1]_{q_{n_1}}x)^{k_1}}{\gamma_{\mu_1,q_{n_1}}(k_1)} \frac{([n_2]_{q_{n_2}}y)^{k_2}}{\gamma_{\mu_2,q_{n_2}}(k_2)}$$

$$\times q_{n_1}^{\frac{k_1(k_1-1)}{2}} q_{n_2}^{\frac{k_2(k_2-1)}{2}} \int_R \int f\left(\frac{n_1 x + \alpha_1}{n_1+\beta_1}, \frac{n_2 y + \alpha_2}{n_2+\beta_2} \right) d_q y d_q x \qquad (5.1)$$

where

$$R = \left[\frac{[k_1+2\mu_1\theta_{k_1}]_{q_{n_1}}}{q_{n_1}^{k_1-2}[n_1]_{q_{n_1}}}, \frac{[k_1+1+2\mu_1\theta_{k_1}]_{q_{n_1}-1}}{q_{n_1}^{k_1-1}[n_1]_{q_{n_1}}} + \frac{1}{[n_1]_{q_{n_1}}} \right]$$

$$\times \left[\frac{[k_2+2\mu_2\theta_{k_2}]_{q_{n_2}}}{q_{n_2}^{k_2-2}[n_2]_{q_{n_2}}}, \frac{[k_2+1+2\mu_2\theta_{k_2}]_{q_{n_2}-1}}{q_{n_2}^{k_2-1}[n_2]_{q_{n_2}}} + \frac{1}{[n_2]_{q_{n_2}}} \right]$$

$$= \left\{ (x,y) \in \mathbb{R}^2 \text{ such that } \frac{[k_1+2\mu_1\theta_{k_1}]_{q_{n_1}}}{q_{n_1}^{k_1-2}[n_1]_{q_{n_1}}} \leq x \leq \frac{[k_1+1+2\mu_1\theta_{k_1}]_{q_{n_1}-1}}{q_{n_1}^{k_1-1}[n_1]_{q_{n_1}}} + \frac{1}{[n_1]_{q_{n_1}}} \text{ and } \right.$$

$$\left. \frac{[k_2+2\mu_2\theta_{k_2}]_{q_{n_2}}}{q_{n_2}^{k_2-2}[n_2]_{q_{n_2}}} \leq y \leq \frac{[k_2+1+2\mu_2\theta_{k_2}]_{q_{n_2}-1}}{q_{n_2}^{k_1-1}[n_2]_{q_{n_2}}} + \frac{1}{[n_2]_{q_{n_2}}} \right\},$$

and

$$E_{\mu_1,q_{n_1}}([n_1]_{q_{n_1}}x) = \sum_{k_1=0}^{\infty} \frac{([n_1]_{q_{n_1}}x)^{k_1}}{\gamma_{\mu_1,q_{n_1}}(k_1)} q_{n_1}^{\frac{k_1(k_1-1)}{2}}, \; E_{\mu_2,q_{n_2}}([n_2]_{q_{n_2}}y) = \sum_{k_2=0}^{\infty} \frac{([n_2]_{q_{n_2}}y)^{k_2}}{\gamma_{\mu_2,q_{n_2}}(k_2)} q_{n_2}^{\frac{k_2(k_2-1)}{2}}.$$

Lemma 5.1. *Let $e_{i,j} : \mathbb{R}_+^2 \to [0,\infty)$ be such that $e_{i,j} = u^i v^j$, $i,j = 0,1,2$ be the two dimensional test functions. Then the q-bivariate operators defined in (5.1) satisfy the following:*

(1) $K^*_{n_1,n_2}(e_{0,0};q_{n_1},q_{n_2};x,y) = 1$

(2) $K^*_{n_1,n_2}(e_{1,0};q_{n_1},q_{n_2};x,y) = \frac{n_1}{(n_1+\beta_1)} \left(\frac{\alpha_1}{n_1} + \frac{1}{[2]_{q_{n_1}}[n_1]_{q_{n_1}}} \right) + \frac{2n_1 q_{n_1}}{[2]_{q_{n_1}}(n_1+\beta_1)} x$

(3) $K^*_{n_1,n_2}(e_{0,1};q_{n_1},q_{n_2};x,y) = \frac{n_2}{(n_2+\beta_2)} \left(\frac{\alpha_2}{n_2} + \frac{1}{[2]_{q_{n_2}}[n_2]_{q_{n_2}}} \right) + \frac{2n_2 q_{n_2}}{[2]_{q_{n_2}}(n_2+\beta_2)} y$

(4) $K^*_{n_1,n_2}(e_{2,0};q_{n_1},q_{n_2};x,y) \le \frac{n_1^2}{(n_1+\beta_1)^2}\left(\frac{\alpha_1^2}{n_1^2}+\frac{2\alpha_1}{[2]_{q_{n_1}}n_1[n_1]_{q_{n_1}}}+\frac{1}{[3]_{q_{n_1}}[n_1]^2_{q_{n_1}}(n_1+\beta_1)}\right)+$
$\frac{n_1^2}{(n_1+\beta_1)^2}\left(\frac{3[1+2\mu_1]_{q_{n_1}}}{[3]_{q_{n_1}}[n_1]_{q_{n_1}}}+\frac{3q_{n_1}}{[3]_{q_{n_1}}[n_1]_{q_{n_1}}}+\frac{4q_{n_1}\alpha_1}{[2]_{q_{n_1}}n_1}\right)x+\frac{3n_1^2}{[3]_{q_{n_1}}(n_1+\beta_1)^2}x^2$

(5) $K^*_{n_1,n_2}(e_{0,2};q_{n_1},q_{n_2};x,y) \le \frac{n_2^2}{(n_2+\beta_2)^2}\left(\frac{\alpha_2^2}{n_2^2}+\frac{2\alpha_2}{[2]_{q_{n_2}}n_2[n_2]_{q_{n_2}}}+\frac{1}{[3]_{q_{n_2}}[n_2]^2_{q_{n_2}}(n_2+\beta_2)}\right)+$
$\frac{n_2^2}{(n_2+\beta_2)^2}\left(\frac{3[1+2\mu_2]_{q_{n_2}}}{[3]_{q_{n_2}}[n_2]_{q_{n_2}}}+\frac{3q_{n_2}}{[3]_{q_{n_2}}[n_2]_{q_{n_2}}}+\frac{4q_{n_2}\alpha_2}{[2]_{q_{n_2}}n_2}\right)y+\frac{3n_2^2}{[3]_{q_{n_2}}(n_2+\beta_2)^2}y^2.$

Lemma 5.2. *The q-bivariate operators defined in (5.1) satisfy the following:*

(1) $K^*_{n_1,n_2}(e_{1,0}-x;q_{n_1},q_{n_2};x,y) = \frac{n_1}{(n_1+\beta_1)}\left(\frac{\alpha_1}{n_1}+\frac{1}{[2]_{q_{n_1}}[n_1]_{q_{n_1}}}\right)+\left(\frac{2n_1q_{n_1}}{[2]_{q_{n_1}}(n_1+\beta_1)}-1\right)x$

(2) $K^*_{n_1,n_2}(e_{0,1}-y;q_{n_1},q_{n_2};x,y) = \frac{n_2}{(n_2+\beta_2)}\left(\frac{\alpha_2}{n_2}+\frac{1}{[2]_{q_{n_2}}[n_2]_{q_{n_2}}}\right)+\left(\frac{2n_2q_{n_2}}{[2]_{q_{n_2}}(n_2+\beta_2)}-1\right)y$

(3) $K^*_{n_1,n_2}\left((e_{1,0}-x)^2;q_{n_1},q_{n_2};x,y\right)$
$\le \frac{n_1^2}{(n_1+\beta_1)^2}\left(\frac{\alpha_1^2}{n_1^2}+\frac{2\alpha_1}{[2]_{q_{n_1}}n_1[n_1]_{q_{n_1}}}+\frac{1}{[3]_{q_{n_1}}[n_1]^2_{q_{n_1}}(n_1+\beta_1)}\right)$
$+\left\{\frac{n_1^2}{(n_1+\beta_1)^2}\left(\frac{3[1+2\mu_1]_{q_{n_1}}}{[3]_{q_{n_1}}[n_1]_{q_{n_1}}}+\frac{3q_{n_1}}{[3]_{q_{n_1}}[n_1]_{q_{n_1}}}+\frac{4q_{n_1}\alpha_1}{[2]_{q_{n_1}}n_1}\right)-\frac{2n_1}{(n_1+\beta_1)}\left(\frac{\alpha_1}{n_1}+\frac{1}{[2]_{q_{n_1}}[n_1]_{q_{n_1}}}\right)\right\}x+$
$\left(\frac{3n_1^2}{[3]_{q_{n_1}}(n_1+\beta_1)^2}-\frac{2n_1q_{n_1}}{[2]_{q_{n_1}}(n_1+\beta_1)}+1\right)x^2$

(4) $K^*_{n_1,n_2}\left((e_{0,1}-y)^2;q_{n_1},q_{n_2};x,y\right)$
$\le \frac{n_2^2}{(n_2+\beta_2)^2}\left(\frac{\alpha_2^2}{n_2^2}+\frac{2\alpha_2}{[2]_{q_{n_2}}n_2[n_2]_{q_{n_2}}}+\frac{1}{[3]_{q_{n_2}}[n_2]^2_{q_{n_2}}(n_2+\beta_2)}\right)$
$+\left\{\frac{n_2^2}{(n_2+\beta_2)^2}\left(\frac{3[1+2\mu_2]_{q_{n_2}}}{[3]_{q_{n_2}}[n_2]_{q_{n_2}}}+\frac{3q_{n_2}}{[3]_{q_{n_2}}[n_2]_{q_{n_2}}}+\frac{4q_{n_2}\alpha_2}{[2]_{q_{n_2}}n_2}\right)-\frac{2n_2}{(n_2+\beta_2)}\left(\frac{\alpha_2}{n_2}+\frac{1}{[2]_{q_{n_2}}[n_2]_{q_{n_2}}}\right)\right\}y+$
$\left(\frac{3n_2^2}{[3]_{q_{n_2}}(n_2+\beta_2)^2}-\frac{2n_2q_{n_2}}{[2]_{q_{n_2}}(n_2+\beta_2)}+1\right)y^2.$

In order to obtain the convergence results for the operators $K^*_{n_1,n_2}(f;q_{n_1},q_{n_2};x,y)$, we take $q=q_{n_1},\ q_{n_2}$ where $q_{n_1},q_{n_2}\in(0,1)$ and satisfy

$$\lim_{n_1,n_2} q_{n_1},q_{n_2}\to 1. \tag{5.2}$$

The modulus of continuity for bivariate case is defined as follows:

For $f\in H_\omega(\mathbb{R}^2_+)$
$\widetilde{\omega}(f;\delta_1,\delta_2)$

$$=\sup_{u,x\ge 0}\left\{\left|f(u,v)-f(x,y)\right|;\ \left|u-x\right|\le\delta_1,\left|v-y\right|\le\delta_2,\ (u,v)\in\mathbb{R}^2_+,\ (x,y)\in\mathbb{R}^2_+\right\}. \tag{5.3}$$

where $H_\omega(\mathbb{R}^+)$ is the space of all real-valued continuous functions f.
Then for all $f\in H_\omega(\mathbb{R}_+)$ $\widetilde{\omega}(f;\delta_1,\delta_2)$ we have the following conditions:

(i) $\lim_{\delta_1,\delta_2\to 0}\widetilde{\omega}(f;\delta_1,\delta_2)\to 0$

(ii) $\mid f(u,v)-f(x,y)\mid\le\widetilde{\omega}(f;\delta_1,\delta_2)\left(\frac{|u-x|}{\delta_1}+1\right)\left(\frac{|v-y|}{\delta_2}+1\right).$

Theorem 5.3. *Let $q=q_{n_1},q_{n_2}$ satisfy (5.2) and $K^*_{n_1,n_2}(f;q_{n_1},q_{n_2},x,y)$ be the operators defined by (5.1). Then for any function $f\in\tilde{C}([0,\infty)\times[0,\infty))$, for $(x,y)\in[0,\infty),\ 0<q_{n_1},q_{n_2}<1$, we have*

$$| K^*_{n_1,n_2}(f;q_{n_1},q_{n_2},x,y) - f(x,y) |$$

$$\leq \omega\left(f;\frac{1}{\sqrt{[n_1]_{q_{n_1}}}},\frac{1}{\sqrt{[n_2]_{q_{n_2}}}}\right)\lambda_{n_1}(x)\lambda_{n_2}(y),$$

where $\lambda_{n_1}(x) = K^*_{n_1,n_2}\left((e_{1,0}-x)^2;q_{n_1},q_{n_2};x,y\right)$, $\lambda_{n_2}(y) = K^*_{n_1,n_2}\left((e_{0,1}-y)^2;q_{n_1},q_{n_2};x,y\right)$ *are defined in Lemma 5.2 and* $\tilde{C}[0,\infty)$ *is the space of uniformly continuous functions on* $\mathbb{R}^+$. *Moreover,* $\widetilde{\omega}(f,\delta_{n_1},\delta_{n_2})$ *is the modulus of continuity of the function* $f \in \tilde{C}\left([0,\infty)\times[0,\infty)\right)$ *defined in (5.3).*

Proof. We can prove easily by using the Cauchy-Schwarz inequality and choosing $\delta_1 = \delta_{n_1} = \sqrt{\frac{1}{[n_1]_{q_{n_1}}}}$ and $\delta_2 = \delta_{n_2} = \sqrt{\frac{1}{[n_2]_{q_{n_2}}}}$. So we omit the details. □

Now we give the rate of convergence of the operators $K^*_{n_1,n_2}(f;q_{n_1},q_{n_2};x,y)$ defined in (5.1) in terms of the elements of the usual Lipschitz class $Lip_M(\nu_1,\nu_2)$.

Let $f \in C([0,\infty)\times[0,\infty))$, $M>0$ and $0<\nu_1,\nu_2\leq 1$. The class $Lip_M(\nu_1,\nu_2)$ is defined by

$$Lip_M(\nu_1,\nu_2) = \{f :| f(u,v)-f(x,y) |\leq M \mid u-x\mid^{\nu_1} \mid v-y\mid^{\nu_2} \ (u,v) \text{ and } (x,y)\in[0,\infty)\}. \tag{5.4}$$

Theorem 5.4. *Let* $K^*_{n_1,n_2}(f;q_{n_1},q_{n_2};x,y)$ *be the operator defined in (5.1). Then for each* $f \in Lip_M(\nu_1,\nu_2)$, $(M>0,\ 0<\nu_1,\nu_2\leq 1)$ *satisfying (5.4) we have*

$$| K^*_{n_1,n_2}(f;q_{n_1},q_{n_2};x,y) - f(x,y) |\leq M\left(\lambda_{n_1}(x)\right)^{\frac{\nu_1}{2}}\left(\lambda_{n_2}(y)\right)^{\frac{\nu_2}{2}},$$

where $\lambda_{n_1}(x)$ *and* $\lambda_{n_2}(y)$ *are defined in Theorem 5.3.*

Proof. We can prove easily it by using (5.4) and Hölder inequality. So we omit the details. □

Acknowledgement. Second author (MN) acknowledges the financial support of University Grants Commission (Govt. of Ind.) for awarding BSR (Basic Scientific Research) Fellowship.

References

[1] A. Aral, V. Gupta, R.P. Agarwal, Applications of q-Calculus in Operator Theory, Springer, New York 2013.

[2] S.N. Bernstein, Démonstration du théoréme de Weierstrass fondée sur le calcul des probabilités, Commun. Soc. Math. Kharkow, 2(13) (2012), 1–2.

[3] B. Cheikh, Y. Gaied, M. Zaghouani, A q-Dunkl-classical q-Hermite type polynomials, Georgian Math. J., 21(2) (2014), 125–137.

[4] A. Ciupa, A class of integral Favard-Szász type operators, Stud. Univ. Babeş-Bolyai, Math., 40(1) (1995), 39–47.

[5] A. Francesco, Korovkin-type Theorems and Approximation by Positive Linear Operators, arXiv:1009.2601v1 [math.CA] 14 Sep 2010.

[6] A.D. Gadzhiev, Theorems of the type of P. P. Korovkin's theorems (Russian), presented at the international conference on the theory of approximation of functions (Kaluga, 1975). Mat. Zametki 20(5) (1976), 781–786.

[7] V. Gupta, C. Radu, Statistical approximation properties of q-Baskokov-Kantorovich operators, Cent. Eur. J. Math., 7 (4) (2009), 809–818.

[8] G. İçōz, B. Çekim, Stancu type generalization of Dunkl analogue of Szász-Kamtrovich operators, Math. Meth. Appl. Sci., (2015) DOI: 10.1002/mma.3602.
[9] G. İçōz, B. Çekim, Dunkl generalization of Szász operators via q-calculus, Jour. Ineq. Appl., 2015 (2015): 284.
[10] L.V. Kantorovich, Sur certains développements suivant les polynômes de la forme de S. Bernstein, I, II, C. R. Acad. URSS, Vol. 563–568 (1930), 595–600.
[11] P. P. Korovkin, Convergence of linear positive operators in the spaces of continuous functions (Russian), Doklady Akad. Nauk. SSSR (N.S.), 90(1953), 961–964.
[12] A. Lupaş, A q-analogue of the Bernstein operator, In Seminar on Numerical and Statistical Calculus, University of Cluj-Napoca, Cluj-Napoca 9 (1987), 85–92.
[13] N.I. Mahmudov, Statistical approximation of Baskakov and Baskakov-Kantorovich operators based on the q-integers, Cent. Eur. Jour. Math., 8(4) (2010), 816–826.
[14] N.I. Mahmudov, V. Gupta, On certain q-analogue of Szász Kantorovich operators, J. Appl. Math. Comput., 37 (2011), 407–419.
[15] M. Mursaleen, A. Khan, Statistical approximation properties of modified q- Stancu-Beta operators, Bull. Malays. Math. Sci. Soc. (2), 36(3) (2013), 683–690.
[16] M. Mursaleen, A. Khan, Generalized q-Bernstein-Schurer operators and some approximation theorems, Jour. Function Spaces Appl., Volume (2013), Article ID 719834, 7 pages.
[17] M. Mursaleen and Md Nasiruzzaman, A Dunkl generalization of q-parametric Száasz-Mirakjan operators, arXiv:1511.06628v1 [math.CA] 19 Nov 2015.
[18] M. Mursaleen, K. J. Ansari and A. Khan, On (p, q)-analogue of Bernstein operators, Appl. Math. Comput., 266 (2015), 874–882 [Erratum: Appl. Math. Comput**.,** 278 (2016) 70–71].
[19] M. Mursaleen, Md. Nasiruzzaman, Some approximation results of q-Szász-Mirakjan-Kantrovich type operators via Dunkl generalization, arXiv:submit/1484906 [math.FA] 18 Feb (2016)
[20] M. Mursaleen, Khursheed J. Ansari, Asif Khan, Some approximation results by (p, q)-analogue of Bernstein-Stancu operators, Appl. Math. Comput**.,** 264 (2015) 392–402 [Corrigendum: Appl. Math. Comput**,** 269 (2015) 744–746].
[21] M. Örkcü, O. Doğru, Weighted statistical approximation by Kantorovich type q-Szász Mirakjan operators, Appl. Math. Comput., 217 (2011), 7913–7919.
[22] G.M. Phillips, Bernstein polynomials based on the q- integers, Ann. Numer. Math., 4 (1997), 511–518.
[23] M. Rosenblum, Generalized Hermite polynomials and the Bose-like oscillator calculus, Oper. Theory, Adv. Appl., 73 (1994), 369–396.
[24] S. Sucu, Dunkl analogue of Szász operators, Appl. Math. Comput., 244 (2014), 42–48.
[25] O. Szász, Generalization of S. Bernstein's polynomials to the infinite interval, J. Res. Natl. Bur. Stand., 45 (1950), 239–245.